

B_s mixing phase and lepton flavor violation in supersymmetric SU(5)

Jae-hyeon Park*

INFN, Sezione di Padova, via F Marzolo 8, I-35131, Padova, Italy

Masahiro Yamaguchi†

Department of Physics, Tohoku University, Sendai 980-8578, Japan

28 September 2008

Abstract

We inspect consequences of the latest B_s mixing phase measurements on lepton flavor violation in a supersymmetric SU(5) theory. The $\mathcal{O}(1)$ phase, preferring a non-vanishing squark mixing, generically implies $\tau \rightarrow (e + \mu) \gamma$ and $\mu \rightarrow e \gamma$. Depending on the gaugino and the scalar mass parameters as well as $\tan\beta$, the rates turn out to be detectable or even already excessive, if the RR mass insertion of down-type squarks is nonzero. We find that it becomes easy to reconcile B_s mixing phase with lepton flavor violation given: gaugino to scalar squared mass ratio around $1/12$, both LL and RR insertions with decent sizes, and low $\tan\beta$.

In the last few years, two experiments at Tevatron have been accumulating information on the mixing of the B_s -meson. The precision of the mass splitting ΔM_s between the two mass eigenstates composed of B_s and $\bar{B}_s$, by now has reached the level of 0.7% [1], which is comparable to that of the B_d -meson [2]. Despite its high accuracy, ΔM_s is not showing any incompatibility with the Standard Model (SM). This should be regarded as yet another triumph of the model. However, a point to keep in mind at the present moment is that it is not easy to separate an extra contribution within ΔM_s , even if one exists, from that of the SM, due to the large theoretical uncertainty around 30% stemming from errors in the $\Delta B = 2$ hadronic matrix element and the Cabibbo–Kobayashi–Maskawa (CKM) quark mixing matrix [3].

On the other hand, the mixing phase, denoted by ϕ_s , does not suffer from these theoretical uncertainties, and one can make a closer connection between its data and a theory possibly involving new physics [4]. Let us choose the notation ϕ_s to represent what is called $\phi_s^{J/\psi\phi}$ by the Heavy Flavor Averaging Group (HFAG) [2]. In the SM, one has $\phi_s \simeq -2\eta\lambda^2 \simeq -0.04$. On the experimental side, it is still much less precise than $\sin 2\beta$. Nonetheless, ϕ_s is already becoming a useful probe into the flavor sector of an extension of the SM. In particular, one could observe an interesting tendency in both data from DØ [5] and CDF [6], that each result appeared to favor a negative $\mathcal{O}(1)$ value of ϕ_s . This tendency came to stand out after the UTfit collaboration, based on the two experiments, reported that their global fit showed a 3.7σ discrepancy of ϕ_s from its SM value [7]. This deviation, however, has decreased to 2.5σ after they updated their analysis including newly available experimental information from DØ [8]. The latest constrained fit by HFAG shows that [2]

$$\phi_s = -0.76^{+0.37}_{-0.33} \quad \text{or} \quad -2.37^{+0.33}_{-0.37}, \quad (1)$$

which is consistent with the SM at the level of 2.4σ . Still, it is too early to draw a definite conclusion. If the difference solidifies, it should be a clean indication of a new source of CP violation.

A supersymmetric extension of the SM has potential new sources of flavor and/or CP violation in its soft supersymmetry breaking terms. It might be conceivable that one of them is revealing its existence through

*jae-hyeon.park@pd.infn.it

†yama@tuhep.phys.tohoku.ac.jp

the above anomaly. We employ the notation of mass insertion parameters, written in the form of $(\delta_{ij}^d)_{AB}$ with the generation indices $i, j = 1, 2, 3$ and the chiralities $A, B = L, R$. We do not only use their usual definition at the weak scale [9], but also borrow the same notation to specify an off-diagonal element of the soft scalar mass matrix at M_{GUT} , the unification scale [10]. For instance, we define $(\delta_{ij}^d)_{LL} \equiv [m_q^2]_{ij}/\tilde{m}^2$ at M_{GUT} , where $\tilde{m}^2$ is the averaged diagonal entry of m_q^2 , the soft scalar mass matrix of the SU(2) doublet squarks in the basis where the down-type quark Yukawa matrix is diagonal. Being a transition between the second and the third families, $B_s - \overline{B}_s$ mixing is naturally associated with $(\delta_{23}^d)_{AB}$. Among the four possibilities, the LR and the RL mass insertions tend to cause an unacceptable change in $B \rightarrow X_s \gamma$ before they can give an appreciable modification to $B_s - \overline{B}_s$ mixing [11]. Therefore, we focus on LL and RR mixings in what follows.

One could think of a more interesting situation by working with a grand unified theory (GUT). Since a single GUT multiplet contains both (s)quarks and (s)leptons, flavor transitions in the two sectors are related [12, 13, 14, 15, 16, 17, 18, 19, 20]. Then, one immediately arrives at the conclusion that the new source of $b \leftrightarrow s$ transition, needed to account for ϕ_s , generically implies lepton flavor violation (LFV) [21]. We wish to consider this scenario in a model independent fashion taking SU(5) as the unified gauge group. This work has at least two differences from the article just referenced. First, we take into account the running effects of squark masses below M_{GUT} . The diagonal components of the squark and the slepton mass matrices grow in the course of running, and this effect is more important to squarks than to sleptons due to the gluino mass contribution. Because of this difference, the gaugino to scalar mass ratio at M_{GUT} plays an important role in determining relative strengths of the two types of flavor violations, hadronic and leptonic. This finding will be demonstrated later in the results. Second, we inspect additional observables such as $\mu \rightarrow e \gamma$, $S_{CP}^{\phi K}$, and neutron electric dipole moment (EDM). In addition to $\tau \rightarrow \mu \gamma$, $\mu \rightarrow e \gamma$ turns out to be highly sensitive to $(\delta_{23}^d)_{RR}$ thanks to $(\delta_{13}^l)_{RR}$ that is radiatively generated from top Yukawa coupling and CKM mixing [10].

In a related work [10], we present a more detailed study on supersymmetric flavor violation in a SU(5) GUT. Let us recapitulate highlights thereof, relevant to the following discussions. The first topic is the connection of a leptonic process to a squark mixing. We ignore the running effects on slepton mixings from neutrino Yukawa couplings below M_{GUT} . In cases where there are sizeable right-handed down-type squark mixings, they lead to LFV decays dominated by chargino loops. If one has a perfect alignment between the mass eigenstates of quarks and leptons, $(\delta_{ij}^d)_{RR}$ implies the transition of $l_j \rightarrow l_i$. However, this straightforward correspondence may be broken by the inclusion of non-renormalizable terms into the superpotential as a solution to the wrong quark-lepton mass relations of the lighter two families. With the assumption that the cutoff scale of the GUT is two orders of magnitude higher than M_{GUT} , one can nevertheless have

$$B(\tau \rightarrow (e + \mu) \gamma) \propto |(\delta_{13}^l)_{LL}|^2 + |(\delta_{23}^l)_{LL}|^2 \approx |(\delta_{13}^d)_{RR}|^2 + |(\delta_{23}^d)_{RR}|^2 + \mathcal{O}[\cos^2 \beta (\delta_{RR}^d)^2], \quad (2)$$

in terms of insertions at M_{GUT} , exploiting the fact that the breakdown of b - τ alignment is suppressed by $\cos \beta$ [12]. Therefore, non-vanishing $(\delta_{23}^d)_{RR}$ causes either $\tau \rightarrow \mu \gamma$ or $\tau \rightarrow e \gamma$ [10]. A tau decay may be linked also to the left-handed squark mixings. One can reuse (2) except that each chirality index should be flipped to the opposite one. Another difference is that the process amplitude is dominated by a neutralino loop, and thus is much smaller than one from a chargino loop, for a given size of mixing.

An analogous statement can be made regarding $\mu \rightarrow e \gamma$, albeit in a somewhat involved form. It is applicable only to the RR mixings, due to the mechanism by which the decay occurs. The branching fraction has a lower bound such that

$$B(\mu \rightarrow e \gamma) \gtrsim k \times \min\{|(\delta_{13}^l)_{RR}|^2, |(\delta_{23}^l)_{RR}|^2\} \cdot [|(\delta_{13}^d)_{RR}|^2 + |(\delta_{23}^d)_{RR}|^2], \quad (3)$$

with the terms suppressed by $\cos^2 \beta$ omitted. The proportionality constant k can be worked out by calculating the rate from a neutralino loop with triple mass insertions $(\delta_{13}^l)_{RR}(\delta_{33}^l)_{RL}(\delta_{32}^l)_{LL}$. The second factor is at least around $(\delta_{13}^d)_{LL}$ which is supposed to have received radiative corrections from top Yukawa coupling

and CKM mixing [22]. Thus, nonzero $(\delta_{23}^d)_{RR}$ gives rise to $\mu \rightarrow e\gamma$, unless there is a fine-tuning among parameters in the superpotential and the soft supersymmetry breaking sector [10].

The second topic, on hadronic processes, is the competition between squark decoupling and the growth of a Δ parameter, as the diagonal components of the squark mass matrix increase. By Δ , we mean the off-diagonal part of a sfermion mass matrix. Suppose that the δ parameters defined above and the gaugino mass $M_{1/2}$ are fixed at M_{GUT} . Imagine that one can increase m_0 , the common diagonal entries of soft squark mass matrix at M_{GUT} , from the value which make the gluino and the squark masses coincide at the weak scale. This value corresponds to $x \equiv M_{1/2}^2/m_0^2 \approx 0.7$. As m_0 increases, $(\Delta_{ij}^d)_{AB} \equiv m_0^2 \times (\delta_{ij}^d)_{AB}$ grows as well, thereby exerting more and more influence on low energy flavor violation such as B_s mixing. At some point, however, squark loop effects begin to decouple as the squarks become too heavy. For $B_s-\overline{B}_s$ mixing, this is around $x = 1/12$. This gaugino to scalar mass ratio could be regarded as a condition for optimizing the sensitivity of a hadronic process to flavor non-universality at M_{GUT} [10]. The importance of this observation is more pronounced when one tries to compare hadronic and leptonic constraints since the latter is monotonically weakened as m_0 is being raised.

Having briefed the reader on qualitative aspects of flavor physics in a supersymmetric GUT, we proceed to computation. We take the same procedure of numerical analysis as in Ref. [10]. As was already mentioned, we restrict ourselves to LL and RR mixings of down-type squarks. Regarding patterns of the two insertions, we consider three scenarios: the LL scenario, the RR scenario, and the $LL = RR$ scenario. The meaning of each name should be self-explanatory except that we set an LL insertion, unless it is a scanning variable, to a value generated by renormalization group (RG) running from the supersymmetry breaking mediation scale M_* down to M_{GUT} , where M_* is taken to be the reduced Planck scale. We do this for $(\delta_{12}^d)_{LL}$ and $(\delta_{13}^d)_{LL}$ as well as $(\delta_{23}^d)_{LL}$. These boundary conditions are given at M_{GUT} with which we solve one-loop RG equations down to the weak scale. We consider only the gluino loop contributions to a quark sector process. We display the portion of the parameter space permitted by each constraint on the complex plane of a GUT scale mass insertion. As for ϕ_s , we use the 90% confidence level (CL) region from HFAG [2],

$$\phi_s \in [-1.26, -0.13] \cup [-3.00, -1.88]. \quad (4)$$

For concreteness, we assume that there is an exact quark-lepton flavor alignment. Regarding $\tau \rightarrow \mu\gamma$, it is straightforward to translate their bounds presented below to a case with quark-lepton misalignment discussed above—interpret $B(\tau \rightarrow \mu\gamma)$ as $B(\tau \rightarrow (e + \mu)\gamma)$. This prescription is applicable to all the three scenarios considered in this work. As for $\mu \rightarrow e\gamma$, barring accidental cancellations, a contour does not need a modification in the RR and $LL = RR$ scenarios, while we do not have a systematic way to account for a misalignment in the LL scenario. We will elaborate on this later on. In order to demonstrate the role of the gaugino to scalar mass ratio, we fix $M_{1/2} = 180$ GeV, which makes the gluino mass be 500 GeV at the weak scale, and then try two different values of $m_0 = 220$ GeV and 600 GeV, corresponding to the right-handed down-type squark masses of 500 GeV and 750 GeV at the weak scale, respectively. The former m_0 results in a benchmark case often considered in the literature, and the latter m_0 optimizes the sensitivity of neutral meson mixing to δ 's at the GUT scale. We also vary $\tan\beta$ from 5 to 10. Other details can be found in Ref. [10], such as experimental inputs in use and ways to impose them as constraints.

First, let us examine the LL mixing scenario. The region preferred by each process is shown in Figs. 1. Among the four figures, Figs. (a) and (b) are for lower m_0 . For this m_0 , one recognizes that the supersymmetric contribution to $B_s-\overline{B}_s$ mixing is not enough to fit the ϕ_s data even if one allows for an $\mathcal{O}(1)$ mass insertion. The dotted contour lines tell us that a maximal alteration in ϕ_s that can be expected is about 0.1. They reveal that the other experimental constraints are not the primary reason why the LL mixing scenario with lower m_0 is inadequate for making an $\mathcal{O}(1)$ change in ϕ_s . The mixing is simply unable to make an enough difference, due to the dilution of squark mixing by gluino mass contribution in the course of RG running down to the weak scale. In Figs. (c) and (d), one can find gray (cyan) regions that lead to ϕ_s within its 90% CL intervals. They involve an $\mathcal{O}(1)$ mass insertion between the second and the third families

of left-handed squarks. For this value of m_0 , squark mixing given at M_{GUT} is less diluted by the running of diagonal components of the mass matrix. However, the supersymmetric disturbance is not only enhanced in B_s mixing, but also in $B \rightarrow X_s \gamma$. Because of this, the bulk of a gray zone is excluded by $B \rightarrow X_s \gamma$. The disturbance in this decay mode grows with $\tan\beta$ [23, 24]. One can see that the $B \rightarrow X_s \gamma$ constraint is severer in Fig. (d) than in Fig. (c), and that there remains a bigger viable corner for lower $\tan\beta$. Also note that for $\tan\beta = 10$, the phase of $(\delta_{23}^d)_{LL}$, needed to fit ϕ_s , modifies $S_{CP}^{\phi K}$ so that it goes out of its current 2σ range, except in a small part of the gray area. Some parts of the regions favored by ϕ_s , give rise to $\tau \rightarrow \mu\gamma$ and/or $\mu \rightarrow e\gamma$ so much as they can be detected at a super B factory [25, 26] or the MEG experiment [27]. LFV rates in those parts increase with $\tan\beta$ enlarging their discovery chance, although large $\tan\beta$ is disfavored by $B \rightarrow X_s \gamma$ and $S_{CP}^{\phi K}$. Remember that the displayed LFV branching ratios have been calculated under the assumption that the quark and the lepton mass eigenstates are aligned. We will come back to consequences of relaxing this assumption later.

Let us turn to the RR scenario. The plots are presented in Figs. 2. Comparing the first two of Figs. 2 with those of Figs. 1, one can notice that gray (cyan) regions are visible here, unlike the LL scenario. This is due to the LL insertion induced by RG running from M_* down to the weak scale [28]. The presence of $(\delta_{23}^d)_{LL}$ enhances the effect of $(\delta_{23}^d)_{RR}$ on $B_s - \bar{B}_s$ mixing [23, 24, 29]. However, those regions leading to ϕ_s within its 90% CL range, are excluded by the current bounds from $\tau \rightarrow \mu\gamma$ [30] and $\mu \rightarrow e\gamma$ [31], even for $\tan\beta$ as low as 5. It seems to be hard to satisfy both ϕ_s and LFV with an RR insertion with low m_0 . This should be contrasted with the LL scenario where LFV was not a major problem. Given non-vanishing $(\delta_{23}^d)_{RR}(M_{\text{GUT}}) = (\delta_{23}^l)_{LL}^*(M_{\text{GUT}})$, $\tau \rightarrow \mu\gamma$ is dominated by a chargino loop, while for mass insertions with the opposite chiralities, it occurs through a neutralino loop. A chargino loop contributes a larger amplitude per mass insertion than a neutralino loop [32]. Therefore, $\tau \rightarrow \mu\gamma$ acts as a tighter restriction here than in the LL scenario. Note that $\mu \rightarrow e\gamma$ occurs as well. This stems from the nonzero $(\delta_{13}^l)_{RR}$ set as a boundary condition at M_{GUT} . This value is expected from the radiative correction from top Yukawa coupling and CKM mixing. Picking up this insertion in addition to $(\delta_{33}^l)_{RL}$ and $(\delta_{32}^l)_{LL}$, a neutralino loop for $\mu \rightarrow e\gamma$ can be completed, which is enhanced by the factor m_τ/m_μ coming from $(\delta_{33}^d)_{RL}$ [12, 32, 33, 34]. Neutron EDM, denoted by d_n , restricts the imaginary part of $(\delta_{23}^d)_{RR}$ with the aid of $(\delta_{23}^d)_{LL}$ [35]. As a result, d_n is setting a limit to which an RR insertion can satisfy ϕ_s , although it is weaker than LFV. Contrasting Figs. 2 (a) and (b) shows that both LFV and d_n constraints become tighter for higher $\tan\beta$. Next, we switch to a higher value of m_0 . Compared to the upper row with lower m_0 , the cases in Figs. (c) and (d) need a smaller size of mass insertion to give an enough contribution to $B_s - \bar{B}_s$ mixing, to its phase in particular. The reason has been already explained. In contrast, LFV is suppressed because of heavier sleptons. These two changes make it easier to fit ϕ_s with smaller LFV rates. Enhancement of hadronic processes, though, leads to a stricter d_n limit. A region allowed by d_n and ΔM_s around the origin, is separated from the ϕ_s region. Recall that d_n is influenced through the combination of $\text{Im}[(\delta_{23}^d)_{LL}(\delta_{23}^d)_{RR}^*]$. Thus the band obeying d_n can be rotated to overlap the gray region by altering $(\delta_{23}^d)_{LL}$ at M_{GUT} . This can happen if $(\delta_{23}^d)_{LL}$ is initially non-vanishing with a complex phase at M_* . Alternatively, the non-renormalizable terms in the superpotential could alter the insertion while it runs from M_* to M_{GUT} . The presented plots are valid for the phase of $(\delta_{23}^d)_{LL}$ equal to $\arg(-V_{ts}^* V_{tb})$. It is noticeable that $B \rightarrow X_s \gamma$ is not playing a very important role. Its branching ratio is not affected as much as in the LL mixing scenario since the supersymmetric amplitude does not interfere with the SM one. Still, the $B \rightarrow X_s \gamma$ ring should be able to touch the gray region for $\tan\beta$ higher than 10. LFV and d_n are also enhanced for high $\tan\beta$. Therefore, lowering $\tan\beta$ helps satisfy LFV and d_n as well as ϕ_s . The contour lines of ϕ_s and a LFV branching fraction show the correlation between them. Suppose that $\tan\beta = 5$. One can find that the region preferred by ϕ_s involves the $\tau \rightarrow \mu\gamma$ rate in the vicinity of the current upper limit. For example, fitting the central value of ϕ_s in (1) causes $B(\tau \rightarrow \mu\gamma)$ to be around 10^{-7} which is already ruled out by the Belle data [30]. The area still surviving could be explored by current and future experiments. The magnitude of mass insertion accessible with the sensitivity of 10^{-8} , attainable at a super B factory, is depicted by a thin circle inside the current upper bound. The prospect may be brighter

according to Ref. [26], which proposes an upper limit of 2×10^{-9} with 75 ab^{-1} . The gray region is also expected to bring about $\mu \rightarrow e\gamma$ at a rate that can be probed by MEG. A more detailed model independent study on the connection between $(\delta_{23}^d)_{RR}$ and $\mu \rightarrow e\gamma$ has been performed in Ref. [10].

The preceding results are based on the supposition that the quark and the lepton mass eigenstates are aligned to each other. With the following modifications, they can be applied to cases where this alignment is disturbed by the Planck-suppressed non-renormalizable operators incorporated to reproduce masses of the lighter quarks and leptons. Replace $\tau \rightarrow \mu\gamma$ by $\tau \rightarrow (e + \mu)\gamma$, i.e. interpret the branching ratio of the former as that of the latter. Obtain a new thick $\tau \rightarrow (e + \mu)\gamma$ ring by expanding the old thick $\tau \rightarrow \mu\gamma$ ring, in order to encompass the events of $\tau \rightarrow e\gamma$ [36]. For this, multiply the old radius by 1.9. Leave the thin $\tau \rightarrow (e + \mu)\gamma$ (namely former $\tau \rightarrow \mu\gamma$) circles untouched. The $\mu \rightarrow e\gamma$ contours should be kept as they are in the RR scenario, and discarded in the LL scenario. The net consequence of these operations is that the current upper bound from $\tau \rightarrow \mu\gamma$ has been relaxed by the factor of 1.9 and $\mu \rightarrow e\gamma$ has been disconnected from the LL mixing. We come back to the plots of RR scenario in Figs. 2. For lower m_0 , the conflict between LFV and ϕ_s is not very much ameliorated, partly due to the still-strong $\tau \rightarrow (e + \mu)\gamma$ and partly due to $\mu \rightarrow e\gamma$. For higher m_0 , the overlap broadens between the zones preferred by LFV and ϕ_s . One can read off the correlation between ϕ_s and $B(\tau \rightarrow (e + \mu)\gamma)$ from their contour lines. Given a prediction of $B(\tau \rightarrow (e + \mu)\gamma)$ and the same future branching ratio reaches of $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$, say 10^{-8} , the chance of discovering either is minimized when the two modes have equal rates. Even in this worst case, a point on the plot could be probed by LFV if it leads to $B(\tau \rightarrow (e + \mu)\gamma)$ above 2×10^{-8} . Indeed, one can find a substantial part of a gray region with this property. Next, we reinterpret Figs. 1 of the LL scenario. There, the role of LFV was not outstanding already before modification. Now, imposing $\tau \rightarrow (e + \mu)\gamma$ instead of $\tau \rightarrow \mu\gamma$ moves its thick circle outside the visible range. This makes the current LFV data further irrelevant to an $\mathcal{O}(1)$ mixing.

Let us digress a little to remark on large neutrino Yukawa couplings. If right-handed neutrinos are heavy, the weak scale mass insertions of sleptons receive corrections from the neutrino Yukawa couplings while running below M_{GUT} [37]. This contribution makes a shift in the position of a LFV circle on the RR mixing plot. This might improve or worsen the compatibility between LFV and ϕ_s , depending on the direction of the displacement. There are cases with specific conditions where one can easily guess the consequences. Suppose that the scalar masses are universal at M_* and that the right-handed neutrinos are integrated out at a single scale M_R . In this case, $(\delta_{23}^d)_{RR}$, displayed in Figs. 2, is assumed to arise solely from neutrino Yukawa couplings. Then a LFV upper bound shrinks by the factor, $\ln(M_*/M_{\text{GUT}})/\ln(M_*/M_R)$ [10], leaving a less room for RR mixing at M_{GUT} than is shown in Figs. 2. One can also apply this method to a case where there is a large hierarchy among the right-handed neutrino masses, by replacing M_R with the largest eigenvalue of M_N [13].

We examine the last scenario with the condition that $(\delta_{23}^d)_{LL} = (\delta_{23}^d)_{RR}$ at M_{GUT} . The results are shown in Figs. 3. Comparing Figs. 3 (a) and (b) with Figs. 2 (a) and (b), it appears that the conflict between LFV and ϕ_s has been much reduced here. Simultaneous presence of LL and RR mixings supplies a reinforced contribution to the B_s mixing even with a smaller size of each insertion about 0.2. On the other hand, the LFV bounds remain almost the same since the dominant source of each mode is $(\delta_{23}^d)_{RR}$, as should be evident from Figs. 1 and 2. Nonetheless, the LFV data shows a disagreement with ϕ_s for lower m_0 , which grows severer for higher $\tan\beta$. Again, raising m_0 to the optimal point, one can enhance supersymmetric effects on $B_s - \overline{B}_s$ mixing while suppressing LFV. Especially, Fig. 3 (c) shows regions well inside the LFV bounds which lead to ϕ_s in perfect agreement with the latest global fit. Part of those regions can satisfy $S_{CP}^{\phi K}$ and d_n as well. Notice that even though d_n is very sensitive to the product of $(\delta_{23}^d)_{LL}$ and $(\delta_{23}^d)_{RR}$, it is not particularly enhanced relative to that in Figs. 2, where the LL insertion is much smaller than here. This is because d_n is a function of $\text{Im}[(\delta_{23}^d)_{LL}(\delta_{23}^d)_{RR}^*]$ at the weak scale and the contribution to this imaginary part arises only through the RG-generated part of $(\delta_{23}^d)_{LL}$. However, these two insertions both of large sizes can generically disturb d_n to a great extent, once one relaxes the assumption that the phases of the LL and

the RR insertions are aligned. This should be taken into account when one tries to guess a situation with two uncorrelated large insertions. On the other hand, it is always possible to escape from d_n if one is willing to tune the relative phase between $(\delta_{23}^d)_{LL}$ and $(\delta_{23}^d)_{RR}$. One finds that $B(B \rightarrow X_s \gamma)$ appears to prefer the left part of the plane. This is because the SM value of the branching ratio, 3.2×10^{-4} , that we use is smaller than the current central value from data [2]. However, there is an enough possibility for the band to be shifted left or right according to the other contributions from loops involving chargino or charged Higgs. Taking only the (half) width of the band as a criterion for an acceptable size of mass insertion, one could regard regions on both sides of the $B \rightarrow X_s \gamma$ curve as acceptable. An area preferred by ϕ_s gives rise to $B(\tau \rightarrow \mu \gamma)$ around 10^{-8} . The rate of $\mu \rightarrow e \gamma$ expected from the same area is around the sensitivity of MEG. In Fig. (d), we vary $\tan\beta$ up to 10. The LFV bounds become tighter. Nevertheless, there are corners of the gray zones that obey all the constraints. Remember that d_n can be loosened by modifying the relative phase between $(\delta_{23}^d)_{LL}$ and $(\delta_{23}^d)_{RR}$. Obviously, the chance of observing LFV at a future experiment increases with $\tan\beta$. Let us comment on more general cases with quark-lepton flavor misalignment. One can convert each plot to a version for misalignment in the same way as one did in the RR scenario, since the LFV modes are dominated by the RR insertion in this scenario as well. The maximum magnitude of insertion set by $\tau \rightarrow \mu \gamma$ should be multiplied by 1.9. For lower m_0 , most [Fig. (a)] or all [Fig (b)] of the region favored by ϕ_s is still excluded by $\mu \rightarrow e \gamma$ although it is a little weaker than $\tau \rightarrow \mu \gamma$ before the conversion. For higher m_0 , the conversion lifts the barrier of LFV even for $\tan\beta = 10$, thereby relieving the tension between ϕ_s and LFV.

Finally, we come to the summary. We have assessed consequences of the latest ϕ_s data on scalar flavor non-universality at the GUT scale within the framework of supersymmetric SU(5) grand unification. We have taken a model independent approach making use of mass insertion parameters. We have examined three patterns of $(\delta_{23}^d)_{LL}$ and $(\delta_{23}^d)_{RR}$: LL , RR , and $LL = RR$. For reconciling ϕ_s with LFV, it greatly helps to choose the optimal value of the GUT scale gaugino to scalar mass ratio, in all these three scenarios. It appears that the most adequate to fit the current value of ϕ_s is $LL = RR$ among the three scenarios. The rest two might still be able to push ϕ_s into its 90% CL range. The barriers to this purpose in the LL scenario are $B \rightarrow X_s \gamma$ and $S_{CP}^{\phi K}$, but there are cases with low $\tan\beta$ where they leave a corner of the parameter space satisfying ϕ_s . In the RR scenario, the major obstacles are LFV and the neutron EDM. Yet, the former is not totally mutually exclusive with ϕ_s , and the latter can be circumvented by a modification to the LL insertion. The neutron EDM is a potential danger in the $LL = RR$ scenario as well depending on the relative phase of the two insertions. Inclusion of Planck-suppressed non-renormalizable terms for fixing the quark-lepton mass relations, in general, affects a LFV bound. This alteration can be estimated by weakening a $\tau \rightarrow \mu \gamma$ bound to that from $\tau \rightarrow (e + \mu) \gamma$. In the two scenarios involving an RR mixing, this reduces the tension between LFV and ϕ_s for higher m_0 , while $\mu \rightarrow e \gamma$ keeps disfavoring lower m_0 . In all cases, low $\tan\beta$ loosens $B \rightarrow X_s \gamma$, $S_{CP}^{\phi K}$, and d_n as well as LFV, providing for more room to accommodate ϕ_s .

We thank Diego Tonelli for useful comments. JhP acknowledges Research Grants funded jointly by the Italian Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR), by the University of Padova and by the Istituto Nazionale di Fisica Nucleare (INFN) within the *Astroparticle Physics Project* and the FA51 INFN Research Project. This research was supported in part by the European Community Research Training Network UniverseNet under contract MRTN-CT-2006-035863. The work of MY was partially supported by the grants-in-aid from the Ministry of Education, Science, Sports and Culture in Japan, No. 16081202 and No. 17340062.

References

- [1] A. Abulencia *et al.* [CDF Collaboration], Phys. Rev. Lett. **97** (2006) 242003 [hep-ex/0609040].

- [2] E. Barberio *et al.* [Heavy Flavor Averaging Group], arXiv:0808.1297 [hep-ex].
- [3] A. Lenz and U. Nierste, JHEP **0706** (2007) 072 [hep-ph/0612167].
- [4] For a review, see e.g. R. Fleischer, arXiv:0802.2882 [hep-ph].
- [5] V. M. Abazov *et al.* [DØ Collaboration], arXiv:0802.2255 [hep-ex].
- [6] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **100** (2008) 161802 [arXiv:0712.2397 [hep-ex]].
- [7] M. Bona *et al.* [UTfit Collaboration], arXiv:0803.0659 [hep-ph].
- [8] M. Ciuchini, talk at 5th International Workshop on the CKM Unitarity Triangle (CKM2008), Roma, September 9–13, 2008, <http://ckm2008.roma1.infn.it/>.
- [9] L. J. Hall, V. A. Kostelecky and S. Raby, Nucl. Phys. **B267** (1986) 415.
- [10] P. Ko, J.-h. Park and M. Yamaguchi, arXiv:0809.2784 [hep-ph].
- [11] G. L. Kane, P. Ko, H. b. Wang, C. Kolda, J.-h. Park and L. T. Wang, Phys. Rev. **D70** (2004) 035015 [hep-ph/0212092]; Phys. Rev. Lett. **90** (2003) 141803 [hep-ph/0304239]; M. Ciuchini, E. Franco, A. Masiero and L. Silvestrini, Phys. Rev. **D67** (2003) 075016 [Erratum-ibid. **D68** (2003) 079901] [hep-ph/0212397].
- [12] S. Baek, T. Goto, Y. Okada and K. Okumura, Phys. Rev. **D64** (2001) 095001 [hep-ph/0104146].
- [13] J. Hisano and Y. Shimizu, Phys. Lett. **B565** (2003) 183 [hep-ph/0303071].
- [14] M. Ciuchini, A. Masiero, L. Silvestrini, S. K. Vempati and O. Vives, Phys. Rev. Lett. **92** (2004) 071801 [hep-ph/0307191].
- [15] B. Dutta and Y. Mimura, Phys. Rev. Lett. **97** (2006) 241802 [hep-ph/0607147]; arXiv:0805.2988 [hep-ph].
- [16] K. Cheung, S. K. Kang, C. S. Kim and J. Lee, Phys. Lett. **B652** (2007) 319 [hep-ph/0702050].
- [17] M. Ciuchini, A. Masiero, P. Paradisi, L. Silvestrini, S. K. Vempati and O. Vives, Nucl. Phys. **B783** (2007) 112 [hep-ph/0702144].
- [18] F. Borzumati, S. Mishima and T. Yamashita, talk at Flavor in the Era of the LHC: 4th Workshop on the Interplay of Flavor and Collider Physics, Geneva, and 4th International Workshop on the CKM Unitarity Triangle (CKM 2006), 12–16 December 2006 Nagoya, arXiv:0705.2664 [hep-ph].
- [19] T. Goto, Y. Okada, T. Shindou and M. Tanaka, Phys. Rev. **D77** (2008) 095010 [arXiv:0711.2935 [hep-ph]].
- [20] J. Hisano and Y. Shimizu, arXiv:0805.3327 [hep-ph].
- [21] J. K. Parry and H. h. Zhang, Nucl. Phys. **B802** (2008) 63 [arXiv:0710.5443 [hep-ph]].
- [22] R. Barbieri and L. J. Hall, Phys. Lett. **B338** (1994) 212 [hep-ph/9408406].
- [23] F. Gabbiani and A. Masiero, Nucl. Phys. **B322** (1989) 235.
- [24] P. Ko and J.-h. Park, arXiv:0809.0705 [hep-ph].
- [25] S. Hashimoto *et al.*, KEK-REPORT-2004-4.

- [26] M. Bona *et al.*, arXiv:0709.0451 [hep-ex].
- [27] A. Baldini *et al.*, Research Proposal to INFN, September 2002, <http://meg.web.psi.ch/>.
- [28] L. Alvarez-Gaume, M. Claudson and M. B. Wise, Nucl. Phys. **B207** (1982) 96.
- [29] M. Ciuchini and L. Silvestrini, Phys. Rev. Lett. **97** (2006) 021803 [hep-ph/0603114]; M. Endo and S. Mishima, Phys. Lett. **B640** (2006) 205 [hep-ph/0603251].
- [30] K. Hayasaka *et al.* [Belle Collaboration], Phys. Lett. **B666** (2008) 16 [arXiv:0705.0650 [hep-ex]].
- [31] M. L. Brooks *et al.* [MEGA Collaboration], Phys. Rev. Lett. **83** (1999) 1521 [hep-ex/9905013].
- [32] J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. **D53** (1996) 2442 [hep-ph/9510309].
- [33] R. Barbieri, L. J. Hall and A. Strumia, Nucl. Phys. **B445** (1995) 219 [hep-ph/9501334]; J. Hisano and D. Nomura, Phys. Rev. **D59** (1999) 116005 [hep-ph/9810479]; L. Calibbi, A. Faccia, A. Masiero and S. K. Vempati, JHEP **0707** 012 (2007) [hep-ph/0610241].
- [34] P. Paradisi, JHEP **0510** (2005) 006 [hep-ph/0505046].
- [35] J. Hisano and Y. Shimizu, Phys. Rev. **D70** (2004) 093001 [hep-ph/0406091]; J. Hisano, M. Kakizaki, M. Nagai and Y. Shimizu, Phys. Lett. **B604** (2004) 216 [hep-ph/0407169].
- [36] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **96** (2006) 041801 [hep-ex/0508012].
- [37] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.




Figure 1: Constraints on the complex plane of $(\delta_{23}^d)_{LL}$, with $(\delta_{12}^d)_{LL}$ and $(\delta_{13}^d)_{LL}$ generated from RG running between the reduced Planck scale and the GUT scale. For $\tau \rightarrow \mu\gamma$, the thick circle is the current upper bound, and the thin circle is an upper bound from the prospective branching ratio limit, 10^{-8} . For $\mu \rightarrow e\gamma$, the thin circle shows the projected bound on the branching ratio, 10^{-13} . A light gray (yellow) region is allowed by ΔM_s , given 30% uncertainty in the $\Delta B = 2$ matrix element, and a gray (cyan) region is further consistent with ϕ_s . The white curves mark a possible improved constraint from ΔM_s with 8% hadronic uncertainty. Of the two sides of the $S_{CP}^{\phi K}$ curve, the excluded one is indicated by thin short lines.




Figure 2: Constraints on the complex plane of $(\delta_{23}^d)_{RR}$, with $(\delta_{ij}^d)_{LL}$ generated from RG running between the reduced Planck scale and the GUT scale. For $\tau \rightarrow \mu\gamma$, the thick circle is the current upper bound, and the thin circles are, from inside, branching ratios of 10^{-8} , 10^{-7} , 10^{-6} , respectively. For $\mu \rightarrow e\gamma$, the thick circle is the current upper bound, and the thin circles are, from inside, branching ratios of 10^{-13} , 10^{-10} , respectively. A light gray (yellow) region is allowed by ΔM_s , given 30% uncertainty in the $\Delta B = 2$ matrix element, and a gray (cyan) region is further consistent with ϕ_s . The white curves running from top to bottom mark a possible improved constraint from ΔM_s with 8% hadronic uncertainty. Of the two sides of the $S_{CP}^{\phi K}$ curve, the excluded one is indicated by thin short lines.




Figure 3: Constraints on the complex plane of $(\delta^d_{23})_{LL} = (\delta^d_{23})_{RR}$, with $(\delta^d_{12})_{LL}$ and $(\delta^d_{13})_{LL}$ generated from RG running between the reduced Planck scale and the GUT scale. For $\tau \rightarrow \mu\gamma$, the thick circle is the current upper bound, and the thin circles are, from inside, branching ratios of 10^{-8} , 10^{-7} , 10^{-6} , respectively. For $\mu \rightarrow e\gamma$, the thick circle is the current upper bound, and the thin circles are, from inside, branching ratios of 10^{-13} , 10^{-10} , respectively. A light gray (yellow) region is allowed by ΔM_s , given 30% uncertainty in the $\Delta B = 2$ matrix element, and a gray (cyan) region is further consistent with ϕ_s . The white curves mark a possible improved constraint from ΔM_s with 8% hadronic uncertainty. Of the two sides of the $S_{CP}^{\phi K}$ curve, the excluded one is indicated by thin short lines.